



Bundesverband der implantologisch tätigen Zahnärzte in Europa e.V.  
European Association of Dental Implantologists

## Update on short, angulated and diameter-reduced implants

11th European Consensus Conference (EuCC) 2016 in Cologne  
6 February 2016



2016

European Association of Dental  
Implantologists (BDIZ EDI)

An der Esche 2 · 53111 Bonn  
Tel. 0228/935 92 44 · Fax 0228/935 92 46  
office-bonn@bdizedi.org  
www.bdizedi.org





Bundesverband der  
implantologisch  
tätigen Zahnärzte  
in Europa

European  
Association  
of Dental  
Implantologists

## Guidelines 2016

### Update on short, angulated and diameter-reduced implants

#### 11th European Consensus Conference (EuCC) 2016 in Cologne

6 February 2016

Authors: Jörg Neugebauer, PhD, DMD  
Hans-Joachim Nickenig M.Sc., PhD, DMD  
Joachim E. Zöller, PhD, MD, DMD  
Department of Cranio-maxillofacial and Plastic Surgery  
and Interdisciplinary Department for Oral Surgery and Implantology  
Centre for Dentistry and Oral and Maxillofacial Surgery,  
University of Cologne, Germany  
Director: Professor DDr Joachim E. Zöller

Chairman: Dr J. Neugebauer (Germany)  
Protocol: Dr F. Vizethum (Germany)  
Participants: Ch. Berger (Germany)  
Dr W. Bolz (Germany)  
Dr A. Bowen (Spain)  
Professor Dr D. Deporter (Canada)  
Professor DDr R. Ewers (Austria)  
Dr P. Fairbairn (United Kingdom)  
Professor Dr A. Felino (Portugal)  
Dr Th. Fortin (France)  
Dr V. Gowd (India)  
Professor Dr M. Kern (Germany)  
Professor Dr P. Kobler (Croatia)  
Professor Dr V. Konstantinovic (Serbia)  
Professor Dr M. Marincola (Italy)  
Dr H.J. Nickenig (Germany)  
Professor Dr H. Özyuvaci (Turkey)  
Professor Dr N. Schmedtmann (Germany)  
Professor DDr J.E. Zöller (Germany)

## 1. Methods

### 1.1 Objective

The purpose of these guidelines is to offer recommendations for clinicians engaging in implant dentistry, enabling them to correctly assess potential indications (and any limitations thereof) for short, angulated or diameter-reduced implants.

BDIZ EDI  
An der Esche 2  
D-53111 Bonn  
GERMANY  
Fon: +49-228-93592-44  
Fax: +49-228-93592-46  
office-bonn@bdizedi.org  
www.bdizedi.org



## 1.2 Introduction

This consensus paper covers only titanium implants typically placed in accordance with the indications recommended by the European Consensus Conference (EuCC, Germany, 6 February 2016).

All consensus recommendations in this paper should be considered as guidelines only. The patient's specific situation is always an important consideration and may justify a deviation from the recommendations of this consensus paper.

## 1.3 Background

Avoiding bone augmentation through reduced-dimension implants and optimum utilization of available bone volume is often recommended being a minimally invasive treatment option <sup>[45]</sup>. To ensure an acceptable treatment outcome, dimension and insertion type must be considered in addition to the number of implants.

## 1.4 Literature search

The Cochrane Library, EMBASE, DIMDI and Medline literature databases were used to conduct a systematic search of recently published data on the use of short, angled or diameter-reduced implants. Selective search criteria were used, including terms such as "short implants", "angulated implants", "angled implants", "tilted implants", "outcome grafting procedure", and "implant -failure". The publications identified by the search were screened by reading their abstracts, and those irrelevant to the subject were identified and excluded. Publications found to be potentially relevant were obtained in full-text form. Multiple review papers with meta-analyses and randomized controlled trials (RCTs), and other prospective and retrospective systematic clinical studies were available on the subject.

## 1.5 Procedure for developing the Consensus Conference guidelines

A preliminary version of this document on which the EuCC based its deliberations was prepared by *Dr J. Neugebauer* of the Interdisciplinary Policlinic for Oral Surgery and Implantology and Department of Oral and Maxillofacial Plastic Surgery at the University of Cologne/Germany. The preliminary report was then reviewed and discussed by the sitting committee members in five steps as follows:

- Reviewing the preliminary draft
- Collecting alternative proposals
- Voting on recommendations and levels of recommendation
- Discussing non-consensual issues
- Final voting

The full text of all (potentially) relevant citations was obtained if necessary and reviewed. Numerous reviews, but few RCTs (randomised controlled trials) or other systematic clinical trials are available on this topic.

## 2. Problem

The application of standard implants in patients with atrophy of their alveolar ridges or large pneumatization of the maxillary sinus cavity often requires the use of hard tissue augmentation procedures <sup>[18, 17]</sup>. These procedures are established, and widely used with success. But depending on level of training of the user and the patient-specific risk factors, complications may occur and affect the postoperative quality of life <sup>[1, 9, 19, 18, 17, 33]</sup>.



### 3. Use of short implants

#### 3.1 Introduction

Short implants are increasingly being discussed as a treatment alternative in situations characterized by limited vertical bone height <sup>[3]</sup>.

Compared to the use of standard implants due to biomechanical considerations (e.g. crown-to-implant ratio, C/R) with short implants may lead to unfavourable loading conditions and complications, including excessive crestal bone loss and implant failure <sup>[23]</sup>. Improvements in implant design and surface along with the use of modified implant insertion methods all are intended to minimize these risks <sup>[14]</sup>.

#### 3.2 Definition of short implants

Implants are usually referred to as short if their designed intrabony length measures  $\leq 8$  mm with diameters  $\geq 3.75$  mm. Standard implants are considered to be those with lengths  $> 8$  mm and diameters  $\geq 3.75$  mm <sup>[43, 47]</sup>. "Ultra-short" implants are considered to be those with lengths less than 6 mm <sup>[13]</sup>.

#### 3.3 Indications for short implants

Short implants are primarily used to avoid bone augmentation procedures in the maxillary and mandibular posterior segments of partially edentulous patients. They are applicable if vertical bone volume is limited by anatomical structures (maxillary sinus, mandibular canal), but there is sufficient alveolar ridge width to permit successful use of implant diameters  $\geq 3.75$  mm. They are also used to support removable overdentures as single or multiple tooth replacements in the anterior jaws <sup>[47, 48]</sup>.

#### 3.4 Current observations

For ultra-short implants, there is insufficient evidence to make recommendations at this time. A review paper from 2015 summarized findings with RCTs on sinus floor elevation with standard length implants or short implants on their own. Five studies reported 16–18 months survival rates for long implants in combination with sinus elevation of 99.5 % (95 % CI: 97.6–99.98 %) and for short implants alone of 99.0 % (95 % CI: 96.4–99.8 %). For shorter observation periods of 8–9 months in three studies, survival rates for long implants were 100 % (95% CI: 97.1–100 %) and for short implants alone 98.2 % (95 % CI: 93.9–99.7 %) <sup>[53]</sup>. These results are supported by other RCTs <sup>[49, 52]</sup>.

The number of RCTs on the use in the mandible is limited <sup>[42]</sup>. In these RCTs, no relevant differences in biological parameters between the use of short and long implants in the posterior mandible were found <sup>[20, 27]</sup>. One group has presented five-year results showing no significant difference for the application of short implants alone as compared to standard implants and vertical augmentation in the mandible <sup>[21]</sup>.

A retrospective comparative analysis also showed no differences between short and long implants for an observation period of five years <sup>[24]</sup>. Meta-analysis showed high survival rates for short implants with moderately rough surfaces <sup>[37]</sup>. Long-term data for observation periods of 10 years for the posterior mandible of partially edentulous patients and 20 years for mandibular overdentures showed favourable results for short, sintered porous-surfaced implants <sup>[15, 16]</sup>.

The literature does show, however, that short implants with a reduced diameter have failure rates of up to 10 % after three to five years <sup>[11]</sup>.



### 3.5 Prevention of complications

Some authors have offered recommendations on how to avoid complications that are mainly biomechanical in nature. These recommendations include:

- Machine-surfaced, short implants should not be used <sup>[37]</sup>
- Short implants should only be used if bone-quality is favourable <sup>[48]</sup>
- Restoration with single crowns <sup>[2, 28, 38, 53]</sup>
- Primary splinting of threaded short implants <sup>[39]</sup>
- Guiding surfaces for lateral movements should be avoided <sup>[10]</sup>
- Insertion at or below bone level with tapered abutment design <sup>[30, 34]</sup>
- The implant surgeon and restorative dentist should have adequate training <sup>[53]</sup>
- For short implants no data available for immediate loading procedures

## 4. Use of angulated implants

### 4.1 Introduction

Angulated standard implant designs or non-angulated ones placed in off-axis (tilted) positions are increasingly being used for the splinted reconstructions of edentulous jaws, again as an alternative treatment option to avoid hard tissue augmentation procedures, but also to increase primary stability for immediate loading procedures with longer implants <sup>[10]</sup>. The objective of having implants in a tilted position is to utilize as much bone as possible, while still avoiding vital adjacent structures (e.g. the mental foramen in the mandible or the maxillary sinus in the maxilla). They also increase the surface area for restorative support (through diverging implant axes) <sup>[4]</sup>. Restorations can be inserted on these implants via angulated abutments.

### 4.2 Current observations

Studies of immediate loading concepts with angulated implants used to support full-arch reconstructions on four or six implants in the maxilla and mandible have provided five-to-ten-year data <sup>[6, 7, 22, 26, 32, 35]</sup>. Favourable survival rates were found following the use of primary splinting of angulated/tilted implants with fixed dental prostheses (FDP) for follow-up intervals of up to 6.5 years <sup>[36]</sup>. Various meta-analyses show no differences compared to conventional implant placement/loading in either survival rates or crestal bone loss in the restoration of atrophied edentulous jaws with FDP and angulated implants after a short and medium observation time <sup>[5, 10, 12, 40]</sup>.

### 4.3 Restoration-related experiences

Using a cantilevered, shortened dental arch with a lack of posterior support resulted in no increased prevalence of oro-mandibular malfunctions <sup>[46]</sup>.

### 4.4 Prevention of complications

- Placement of angulated and immediately restored implants should be effected with sufficient primary stability (and splinted with immediate restoration)
- For anatomically and prosthetically correct angulated implant placement, a pre-operative 3D computer-based diagnosis is recommended
- The implant surgeon and restorative dentist must have adequate training



## 5. Use of diameter-reduced implants

### 5.1 Definition

Diameter-reduced implants (DRIs) can be defined as those with intraosseous diameters below 3.5 mm for placement in sites with reduced alveolar ridge bone width <sup>[47]</sup>. Implants with diameters less than 2.7 mm are referred to as mini-implants <sup>[25, 50]</sup>.

### 5.2 Current observations

Diameter-reduced implants generally have high survival rates (> 90 %), assuming careful patient selection, assessment of bone density, the clinical approach and the experience of the user <sup>[29, 31, 47, 51]</sup>. DRIs are also applicable in the posterior region with high success rates <sup>[31]</sup>.

Many retrospective studies are available for mini-implants. However, meta-analyses with prospective and/or randomized trials show only short-term results and/or increased failure rates <sup>[8, 31, 44]</sup>. Furthermore, in a recent literature review, it was determined that short mini-implants ( $\leq 13$  mm) will be lost more frequently under load than longer ones ( $> 13$  mm) <sup>[51]</sup>.

### 5.3 Prevention of complications

- Mini-implants have an increased risk of implant loss
- Short mini-implants should be avoided <sup>[54]</sup>

## 6. Recommendations for short, angulated or diameter-reduced implants

Provided the specific treatment parameters are observed, the use of short, angulated or diameter-reduced implants in sites with reduced bone volume can be a reliable treatment option, given the risks associated with the use of standard-dimension implants in combination with augmentation procedures. The implant surgeon and the restorative dentist must have appropriate training to choose the best possible therapy for each patient <sup>[41]</sup>.

Cologne, 6 February 2016

Professor Dr. Joachim E. Zöller  
Vice President

Dr. Jörg Neugebauer  
Chairman of EuCC



## References

1. Aghaloo TL, Moy PK. Which hard tissue augmentation techniques are the most successful in furnishing bony support for implant placement? *Int J Oral Maxillofac Implants* 2007; 22 Suppl: 49-70.
2. Al-Ansari A. Short implants supporting single crowns in atrophic jaws. *Evid Based Dent* 2014; 15: 85-86.
3. Aloy-Prosper A, Penarrocha-Oltra D, Penarrocha-Diago M, Penarrocha-Diago M. The outcome of intraoral onlay block bone grafts on alveolar ridge augmentations: a systematic review. *Med Oral Patol Oral Cir Bucal* 2015; 20: e251-258.
4. Aparicio C, Perales P, Rangert B. Tilted implants as an alternative to maxillary sinus grafting: a clinical, radiologic, and periosteal study. *Clin Implant Dent Relat Res* 2001; 3: 39-49.
5. Asawa N, Bulbule N, Kakade D, Shah R. Angulated implants: an alternative to bone augmentation and sinus lift procedure: systematic review. *J Clin Diagn Res* 2015; 9: ZE10-13.
6. Babbush CA, Kanawati A, Brokloff J. A new approach to the All-on-Four treatment concept using narrow platform NobelActive implants. *The Journal of oral implantology* 2013; 39: 314-325.
7. Balshi TJ, Wolfinger GJ, Schlauch RW, Balshi SF. A Retrospective Analysis of 800 Branemark System Implants Following the All-on-Four Protocol. *Journal of prosthodontics : official journal of the American College of Prosthodontists* 2014; 23: 83-88.
8. Bidra AS, Almas K. Mini implants for definitive prosthodontic treatment: a systematic review. *The Journal of prosthetic dentistry* 2013; 109: 156-164.
9. Chiapasco M, Casentini P, Zaniboni M. Bone augmentation procedures in implant dentistry. *Int J Oral Maxillofac Implants* 2009; 24 Suppl: 237-259.
10. Chrcanovic BR, Albrektsson T, Wennerberg A. Tilted versus axially placed dental implants: a meta-analysis. *J Dent* 2015; 43: 149-170.
11. das Neves FD, Fones D, Bernardes SR, do Prado CJ, Neto AJ. Short implants—an analysis of longitudinal studies. *Int J Oral Maxillofac Implants* 2006; 21: 86-93.
12. Del Fabbro M, Ceresoli V. The fate of marginal bone around axial vs. tilted implants: a systematic review. *Eur J Oral Implantol* 2014; 7 Suppl 2: S171-189.
13. Deporter D, Ogiso B, Sohn DS, Ruljancich K, Pharoah M. Ultrashort sintered porous-surfaced dental implants used to replace posterior teeth. *J Periodontol* 2008; 79: 1280-1286.
14. Deporter D. Short dental implants: what works and what doesn't? A literature interpretation. *Int J Periodontics Restorative Dent* 2013; 33: 457-464.
15. Deporter D, Pharoah M, Yeh S, Todescan R, Atenafu EG. Performance of titanium alloy sintered porous-surfaced (SPS) implants supporting mandibular overdentures during a 20-year prospective study. *Clin Oral Implants Res* 2014; 25: e189-195.
16. Deporter DA, Kermalli J, Todescan R, Atenafu E. Performance of sintered, porous-surfaced, press-fit implants after 10 years of function in the partially edentulous posterior mandible. *Int J Periodontics Restorative Dent* 2012; 32: 563-570.
17. Esposito M, Grusovin MG, Worthington HV, Coulthard P. Interventions for replacing missing teeth: bone augmentation techniques for dental implant treatment. *Cochrane Database Syst Rev* 2006; CD003607.
18. Esposito M, Grusovin MG, Kwan S, Worthington HV, Coulthard P. Interventions for replacing missing teeth: bone augmentation techniques for dental implant treatment. *Cochrane Database Syst Rev* 2008; CD003607.
19. Esposito M, Felice P, Worthington HV. Interventions for replacing missing teeth: augmentation procedures of the maxillary sinus. *Cochrane Database Syst Rev* 2014; 5: CD008397.
20. Esposito M, Barausse C, Pistilli R, Checchi V, Diazzi M, Gatto MR et al. Posterior jaws rehabilitated with partial prostheses supported by 4.0 x 4.0 mm or by longer implants: Four-month post-loading data from a randomised controlled trial. *Eur J Oral Implantol* 2015; 8: 221-230.
21. Felice P, Cannizzaro G, Barausse C, Pistilli R, Esposito M. Short implants versus longer implants in vertically augmented posterior mandibles: a randomised controlled trial with 5-year after loading follow-up. *Eur J Oral Implantol* 2014; 7: 359-369.
22. Ferreira EJ, Kuabara MR, Gulinelli JL. „All-on-four“ concept and immediate loading for simultaneous rehabilitation of the atrophic maxilla and mandible with conventional and zygomatic implants. *The British journal of oral & maxillofacial surgery* 2010; 48: 218-220.
23. Garaicoa-Pazmino C, Suarez-Lopez del Amo F, Monje A, Catena A, Ortega-Oller I, Galindo-Moreno P et al. Influence of crown/implant ratio on marginal bone loss: a systematic review. *J Periodontol* 2014; 85: 1214-1221.
24. Gentile MA, Chuang SK, Dodson TB. Survival estimates and risk factors for failure with 6 x 5.7-mm implants. *The International journal of oral & maxillofacial implants* 2005; 20: 930-937.
25. Gleiznys A, Skirbutis G, Harb A, Barzdziukaite I, Grinyte I. New approach towards mini dental implants and small-diameter implants: an option for long-term prostheses. *Stomatologija / issued by public institution „Odontologijos studija“ ... [et al.]* 2012; 14: 39-45.
26. Graves S, Mahler BA, Javid B, Armellini D, Jensen OT. Maxillary all-on-four therapy using angled implants: a 16-month clinical study of 1110 implants in 276 jaws. *Dent Clin North Am* 2011; 55: 779-794.
27. Gulje F, Abrahamsson I, Chen S, Stanford C, Zadeh H, Palmer R. Implants of 6 mm vs. 11 mm lengths in the posterior maxilla and mandible: a 1-year multicenter randomized controlled trial. *Clin Oral Implants Res* 2013; 24: 1325-1331.
28. Hasan I, Bourauel C, Mundt T, Heinemann F. Biomechanics and load resistance of short dental implants: a review of the literature. *ISRN Dent* 2013; 2013: 424592.
29. Hasan I, Bourauel C, Mundt T, Stark H, Heinemann F. Biomechanics and load resistance of small-diameter and mini dental implants: a review of literature. *Biomedizinische Technik. Biomedical engineering* 2014; 59: 1-5.



30. Hentschel A, Herrmann J, Glauche I, Vollmer A, Schlegel KA, Lutz R. Survival and patient satisfaction of short implants during the first 2 years of function: a retrospective cohort study with 694 implants in 416 patients. *Clin Oral Implants Res* 2015.
31. Klein MO, Schiegnitz E, Al-Nawas B. Systematic review on success of narrow-diameter dental implants. *Int J Oral Maxillofac Implants* 2014; 29 Suppl: 43-54.
32. Landazuri-Del Barrio RA, Cosyn J, De Paula WN, De Bruyn H, Marcantonio E, Jr. A prospective study on implants installed with flapless-guided surgery using the all-on-four concept in the mandible. *Clin Oral Implants Res* 2013; 24: 428-433.
33. Lee SA, Lee CT, Fu MM, Elmusalati W, Chuang SK. Systematic review and meta-analysis of randomized controlled trials for the management of limited vertical height in the posterior region: short implants (5 to 8 mm) vs longer implants (> 8 mm) in vertically augmented sites. *Int J Oral Maxillofac Implants* 2014; 29: 1085-1097.
34. Lombardo G, Corrocher G, Pighi J, Faccioni F, Rovera A, Marincola M et al. The impact of subcrestal placement on short locking-taper implants placed in posterior maxilla and mandible: a retrospective evaluation on hard and soft tissues stability after 2 years of loading. *Minerva Stomatol* 2014; 63: 391-402.
35. Malo P, Rangert B, Nobre M. „All-on-Four“ immediate-function concept with Branemark System implants for completely edentulous mandibles: a retrospective clinical study. *Clin Implant Dent Relat Res* 2003; 5 Suppl 1: 2-9.
36. Malo P, de Araujo Nobre M, Lopes A, Francischone C, Rigolizzo M. „All-on-4“ immediate-function concept for completely edentulous maxillae: a clinical report on the medium (3 years) and long-term (5 years) outcomes. *Clinical implant dentistry and related research* 2012; 14 Suppl 1: e139-150.
37. Menchero-Cantalejo E, Barona-Dorado C, Cantero-Alvarez M, Fernandez-Caliz F, Martinez-Gonzalez JM. Meta-analysis on the survival of short implants. *Med Oral Patol Oral Cir Bucal* 2011; 16: e546-551.
38. Mezzomo LA, Miller R, Triches D, Alonso F, Shinkai RS. Meta-analysis of single crowns supported by short (<10 mm) implants in the posterior region. *J Clin Periodontol* 2014; 41: 191-213.
39. Misch CE, Steingra J, Barboza E, Misch-Dietsh F, Cianciola LJ, Kazor C. Short dental implants in posterior partial edentulism: a multicenter retrospective 6-year case series study. *J Periodontol* 2006; 77: 1340-1347.
40. Monje A, Chan HL, Suarez F, Galindo-Moreno P, Wang HL. Marginal bone loss around tilted implants in comparison to straight implants: a meta-analysis. *Int J Oral Maxillofac Implants* 2012; 27: 1576-1583.
41. Monteiro DR, Silva EV, Pellizzer EP, Filho OM, Goiato MC. Posterior partially edentulous jaws, planning a rehabilitation with dental implants. *World J Clin Cases* 2015; 3: 65-76.
42. Nisand D, Picard N, Rocchietta I. Short implants compared to implants in vertically augmented bone: a systematic review. *Clin Oral Implants Res* 2015.
43. Olate S, Lyrio MC, de Moraes M, Mazzonetto R, Moreira RW. Influence of diameter and length of implant on early dental implant failure. *Journal of oral and maxillofacial surgery : official journal of the American Association of Oral and Maxillofacial Surgeons* 2010; 68: 414-419.
44. Ortega-Oller I, Suarez F, Galindo-Moreno P, Torrecillas-Martinez L, Monje A, Catena A et al. The Influence of Implant Diameter Upon its Survival: A Meta-Analysis Based on Prospective Clinical Trials. *Journal of periodontology* 2013.
45. Pommer B, Mailath-Pokorny G, Haas R, Busenlechner D, Furhauser R, Watzek G. Patients' preferences towards minimally invasive treatment alternatives for implant rehabilitation of edentulous jaws. *Eur J Oral Implantol* 2014; 7 Suppl 2: S91-109.
46. Reissmann DR, Heydecke G, Schierz O, Marre B, Wolfart S, Strub JR et al. The randomized shortened dental arch study: temporomandibular disorder pain. *Clin Oral Investig* 2014; 18: 2159-2169.
47. Renouard F, Nisand D. Impact of implant length and diameter on survival rates. *Clinical oral implants research* 2006; 17 Suppl 2: 35-51.
48. Romeo E, Bivio A, Mosca D, Scanferla M, Ghisolfi M, Storelli S. The use of short dental implants in clinical practice: literature review. *Minerva stomatologica* 2010; 59: 23-31.
49. Schincaglia GP, Thoma DS, Haas R, Tutak M, Garcia A, Taylor TD et al. Randomized controlled multicenter study comparing short dental implants (6 mm) versus longer dental implants (11-15 mm) in combination with sinus floor elevation procedures. Part 2: clinical and radiographic outcomes at 1 year of loading. *J Clin Periodontol* 2015; 42: 1042-1051.
50. Shatkin TE, Shatkin S, Oppenheimer BD, Oppenheimer AJ. Mini dental implants for long-term fixed and removable prosthetics: a retrospective analysis of 2514 implants placed over a five-year period. *Compend Contin Educ Dent* 2007; 28: 92-99; quiz 100-101.
51. Sohrabi K, Mushantat A, Esfandiari S, Feine J. How successful are small-diameter implants? A literature review. *Clinical oral implants research* 2012; 23: 515-525.
52. Thoma DS, Haas R, Tutak M, Garcia A, Schincaglia GP, Hammerle CH. Randomized controlled multicentre study comparing short dental implants (6 mm) versus longer dental implants (11-15 mm) in combination with sinus floor elevation procedures. Part 1: demographics and patient-reported outcomes at 1 year of loading. *J Clin Periodontol* 2015; 42: 72-80.
53. Thoma DS, Zeltner M, Husler J, Hammerle CH, Jung RE. EAO Supplement Working Group 4 - EAO CC 2015 Short implants versus sinus lifting with longer implants to restore the posterior maxilla: a systematic review. *Clin Oral Implants Res* 2015.
54. Wang HL, Okayasu K, Fu JH, Hamerink HA, Layher MG, Rudek IE. The success rate of narrow body implants used for supporting immediate provisional restorations: a pilot feasibility study. *Implant dentistry* 2012; 21: 467-473.